




Chapter 11: Bifurcation analysis



$$\frac{dR}{dt} = rR(1 - R/K) - \frac{bR^2 N}{h^2 + R^2} \quad \text{and} \quad \frac{dN}{dt} = \frac{cbNR^2}{h^2 + R^2} - dN ,$$


Hopf bifurcation




$$\frac{dR}{dt} = rR(1 - R/K) - \frac{bR^2 N}{h^2 + R^2} \quad \text{and} \quad \frac{dN}{dt} = \frac{cbNR^2}{h^2 + R^2} - dN ,$$

Transcritical bifurcation

(a)




(b)






$$\frac{dR}{dt} = rR\left(1 - \frac{R}{K}\right) - \frac{bR^2N}{h^2 + R^2} \quad \text{and} \quad \frac{dN}{dt} = \frac{cbNR^2d}{h^2 + R^2} - dN ,$$

Saddle node bifurcation




$$\frac{dN}{dt} = \frac{cbNR^2}{h^2 + R^2} - dN - eN^2$$

Pitchfork bifurcation




$$\frac{dN_1}{dt} = i + rN_1(1 - N_1 - cN_2) \quad \text{and} \quad \frac{dN_2}{dt} = i + rN_2(1 - N_2 - cN_1) .$$

Subcritical Hopf and Pitchfork



Chaos in a one consumer two resources model

$$\begin{aligned}\frac{dR_1}{dt} &= R_1(1 - R_1 - \alpha_{12}R_2) - a_1R_1N, \\ \frac{dR_2}{dt} &= R_2(1 - R_2 - \alpha_{21}R_1) - a_2R_2N, \\ \frac{dN}{dt} &= N(ca_1R_1 + ca_2R_2 - 1),\end{aligned}$$



$$a_{12}=1, a_{21}=1.5$$

$a_2=1, c=0.5$:
predator needs R_1


Vary a_1

Chaos in a one consumer two resources model


$$\begin{aligned}\frac{dR_1}{dt} &= R_1(1 - R_1 - \alpha_{12}R_2) - a_1R_1N, \\ \frac{dR_2}{dt} &= R_2(1 - R_2 - \alpha_{21}R_1) - a_2R_2N, \\ \frac{dN}{dt} &= N(ca_1R_1 + ca_2R_2 - 1),\end{aligned}$$

(a)



$a_1=6$





Hopf at $a_1=5.5$



$a_1=8$



$a_1=10$



Period doubling cascade

